

• • •

by Artikel Tnjpr

Submission date: 21-Jun-2022 01:48PM (UTC+0700)

Submission ID: 1860622108

File name: C.1.c.23-TJNPR-2021-M284_Galley_Proof_C.pdf (867.21K)

Word count: 5032

Character count: 28038



Tropical Journal of Natural Product Research

Available online at <https://www.tjnpr.org>

Original Research Article



Phytochemical Constituents and Antimicrobial Activity of *Elaeocarpus sphaericus* Schum Seed Extract

Cicilia N. Primiani^{1*}, Pujiati Pujiati¹, Mohammad A. Setiawan²¹Department of Biology Education, Universitas PGRI Madiun, Madiun, East Java, Indonesia²Department of Chemical Engineering, Universitas PGRI Madiun, Madiun, East Java, Indonesia

ARTICLE INFO

Article history:

Received 25 June 2021

Revised 15 July 2021

Accepted 19 October 2021

Published online 02 November 2021

ABSTRACT

The use of medicinal plants in health care services has tremendously increased. Medicinal plants contain bioactive compounds that are very important in biological activities and may possess antioxidant, anticancer, and antimicrobial properties. Therefore, this study was aimed at identifying the phytochemical constituents of *Elaeocarpus sphaericus* Schum (genitri) seed extract and determining its antibacterial activity. *E. sphaericus* seeds were obtained, dried, and prepared into powder form. The seed powder was extracted with methanol. Phytochemical analysis was conducted on the methanol seed extract using the Liquid Chromatography-Mass Spectrometry (LC-MS) technique. Antibacterial sensitivity testing was carried out with the agar disk diffusion method. The result of the phytochemical screening revealed that *E. sphaericus* seed extract contains 72 compounds which include dicarboxylic acid, aromatic acid, ester, glucose, coumarin, alkaloid, flavonoid, tannin, glycoside, steroid, terpenoid, quinone, and coumestan. Each phytochemical compound varies in composition, with caffeic acid (3.12%) being the highest. The antibacterial sensitivity testing of the 15 and 40% *E. sphaericus* seed extracts against *E. coli* indicated inhibition zone diameters of 7.25 and 7.75 mm, respectively, while values of 7.5 and 9 mm, respectively were recorded for *L. acidophilus*. Also, the antibacterial activity of the seed extract was found to be concentration-dependent. The findings of this study reveal that *E. sphaericus* seed extract contain several phytochemical compounds and has antibacterial activity against *E. coli* and *L. acidophilus*, thereby making it a potential antibacterial agent.

Keywords: Antibacterial, *Elaeocarpus sphaericus* Schum, Metabolites, Phytochemicals.

Copyright: © 2021 Primiani *et al.* This is an open-access article distributed under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Introduction

Biodiversity provides plants that can be used for medicinal purposes. For ages, traditional medical therapies in many countries have used plants to cure a variety of illnesses.¹⁻³ The World Health Organization (WHO) showed that about 75% of the world's population uses herbal extracts for medicine.^{2,4} Furthermore, phytochemical compounds found in plants have the potential to be used as drugs.^{5,6} Several studies have been conducted to investigate their activities in medicinal plants and treatments.^{6,7} Most of the plants that can be used as antioxidants, anticancer, and detoxifying agents, have secondary metabolites such as alkaloids, glycosides, terpene, tannins, steroids, and flavonoids.⁸⁻¹⁰ Meanwhile, the rhizome extract of *Nelumbo nucifera* can be used as an antidiabetic and anti-inflammatory agent because it contains steroidal triterpenoid compounds.⁹ Panduratin A is a phytochemical in the extract of *Boesenbergia rotunda* and it exhibits anti-SARS-CoV-2 activity.¹¹ In essence, various phytochemical compounds have been investigated for treating various infections such as compounds that function as antimicrobials.¹²⁻¹⁴ Genitri (Javanese) holds a taxonomic name from the Elaeocarpaceae family. This plant originated from Java island, and it is currently distributed across Indonesia,¹⁵ with the complete species name

Elaeocarpus sphaericus Schum.

Several previous studies reported that parts of genitri (seeds, fruits, and leaves) have antidiabetic, antiviral, and antimalarial activities.¹⁶⁻¹⁹ There is a dearth of information on the antibacterial activity of seeds of genitri. The seeds contain more complex secondary metabolites than the leaves and fruits and have biologically active phytoconstituents.²⁰⁻²³

E. coli is the most common gram-negative pathogenic bacteria causing a diverse range of clinical diseases that affect all age groups. The pathogen has the potential to invade many tissues and cause infection in any age group.²⁴ Meanwhile, *L. acidophilus* is a gram-positive bacterium which is widely used as a probiotic for humans and animals.^{25,26} Agar disc-diffusion bioassay is widely used in many micro- and nanobiology laboratories. The advantages of the disc diffusion bioassay are ease of use, low cost, capacity to test a large number of microorganisms, and the ability to understand the findings.^{27,28} Furthermore, there are limited reports on the phytochemical composition of the genitri seed extract. The phytochemical studies performed on the Elaeocarpaceae family, particularly on the seed extract were qualitative test.²³ The qualitative test can not reveal the specific phytochemical constituents of the extract. The present study was therefore conducted to perform a phytochemical screening and investigate the antibacterial properties of *Elaeocarpus sphaericus* Schum seed extract.

Materials and Methods

Sample collection

Genitri seeds were picked up from the plants collected from the forest edge area of Poncol Village, Magetan city, East Java, Indonesia, in February 2020. The seeds were identified at the Taxonomy Laboratory of the Universitas PGRI Madiun. They were assigned the following

*Corresponding author. E mail: primiani@unipma.ac.id
Tel: +62 815-5654-1989

Citation: Primiani CN, Pujiati, Setiawan MA. Phytochemical Constituents and Antimicrobial Activities of *Elaeocarpus sphaericus* Schum Seed Extract. Trop J Nat Prod Res. 2021; 5(10):1775-1781. doi.org/10.26538/tjnpr/v5i10.13

Official Journal of Natural Product Research Group, Faculty of Pharmacy, University of Benin, Benin City, Nigeria.

identification number: 0023/Taxo-Plant/Biology/IV/2021. Genitri fruit was drained, peeled, and separated between the flesh and seeds. Then, the genitri seeds were dried by aeration and crushed by pounding. It was filtered through a sieve until a homogeneous powder was obtained.

Preparation of *Elaeocarpus sphaericus* seed extract

Each genitri seed powder was weighed (0.5 – 2 g) and macerated with 95% methanol in a ratio of 1:5 (sample: methanol) for 24 h at cold temperature in a closed bottle. Then, the sample solutions were filtered using an Erlenmeyer vacuum filter to obtain the filtrate and dregs. These steps were repeated three times. Subsequently, the obtained filtrates were mixed and evaporated using a rotary evaporator to separate and obtain the methanol solvent and the semi-viscous extract. Meanwhile, the extract was dissolved using methanol solvent to a concentration below 0.1 mg/mL and homogenized using a vortex tube to obtain a homogeneous solution. To separate the solids, the solution was centrifuged at 8000 rpm for 10 minutes. An aliquot of 2 mL of supernatant was used in the protein precipitation stage. The supernatant was mixed with 3 mL of acetonitrile, acidified with 0.2% formic acid, and centrifuged at 8000 rpm for 30 seconds. Furthermore, the supernatant obtained was then used for analysis.

Fractionation of *Elaeocarpus sphaericus* seed extract

The extract of *Elaeocarpus sphaericus* seeds was purified through solid-phase extraction using C18 Sep-Pak, and the cartridge column C18 was acclimated with a 1 mL solution of 80% acetonitrile. Also, an aliquot of 0.5 mL of the solution was put in a container and 1 mL of the protein precipitate was added into the Sep-Pak column. Then, 0.5 mL of the resulting solution was collected. About 0.25 mL of 200 mM ammonium formate in a 50:50 solution (acetonitrile: methanol) was added into the Sep-Pak column. Then, 0.5 mL of the resulting solution was obtained and 0.2 mL of 25:75 solution (acetonitrile: buffer) was added to 25 mM ammonium formate (pH 4.5). Furthermore, the solution was ready for injection in liquid chromatography-mass spectrometry (LC-MS). The solution was filtered using a 0.45 µm cellulose acetate filter membrane and the degassing procedure was conducted. Also, the sample was injected into the LC-MS system and analyzed.

Antimicrobial sensitivity testing

Escherichia coli and *Lactobacillus acidophilus* were collected from the Microbiology Laboratory, Biology Department, Universitas PGRI Madiun, Indonesia. This study used 40 and 15% concentrations of genitri seed extract. Details for each concentration were 4 g of powdered genitri in 10 mL of water (40%) and 1.5 g of powder of genitri in 10 mL of water (15%). Previous studies showed that these concentrations can be used as an antibacterial test solution.²⁹ In addition, a sheet of filter paper was placed on the surface of the half-solidified nutrient agar (NA) medium and incubated at 37°C for 24 hours. The antibacterial activity was measured by the diameter of the clear zone on NA medium that had been inoculated with *E. coli* or *L. acidophilus*. The procedure involved the addition of 1 mL of bacterial culture from 10⁻⁸ dilution and 1 mL of chloramphenicol solution to NA medium in a petri dish and allowed to solidify. After solidifying, a paper disc that had been soaked in the sample (for two hours) was taken with tweezers and placed on the medium. Each medium contained six paper discs, and the cultures were later incubated upside down for 18-24 hours at 37 °C.³⁰ The size of the antibacterial inhibition zone was determined by measuring the clear zone on a petri dish and subtracting the diameter of the paper disc. These measurements were done by a caliper. The inhibition index was calculated using the following formula.³¹

$$\text{Inhibition index} = \frac{\text{Control clear zone diameter} - \text{Treatment clear zone diameter}}{\text{Disc paper diameter}}$$

From the results of these measurements, the inhibitory effectiveness value can then be calculated based on the following equation:^{32,33}

$$E = \frac{D}{Da} \times 100\%$$

Where E: Inhibitory effectiveness (%); D: Diameter of inhibition zone of plant material (mm); Da: Diameter of antibiotic inhibition zone (mm).


Results and Discussion

The results of the phytochemical screening of genitri seed extract using the LC-MS method are shown in Figure 1 and Table 1. Caffeic acid has the highest concentration in genitri seed extract with a value of 3.12%. LC-MS spectrum analysis identified and classified 72 compounds which include dicarboxylic acid, aromatic acid, ester, glucose, coumarin, alkaloids, flavonoids, tannins, glycosides, steroids, terpenoids, quinones, and coumestans. Secondary metabolites are known to have a specific function as antibacterial agents. This observation is in agreement with a study conducted by Tripathy et al.,²⁰ where they reported that genitri seeds contain secondary metabolites such as alkaloids, phenols, phytosterols, amino acids, flavonoids, and terpenoids. The ethanol/ ether extract of genitri contains secondary metabolites of alkaloids, flavonoids, carbohydrates, proteins, and tannins. In contrast, the aqueous extract contains carbohydrates, proteins, tannins, indoleizidine alkaloids, isoelaecarpine, epiisoelaecarpine, epielaecarpine, alloelaecarpine, and pseudoepiisoelaecarpine.¹⁸ The antibacterial activity of the seed extract was observed due to the presence of several compounds such as flavonoids, saponins, tannins, and alkaloids.^{20,21}

The results (Table 2 and Figure 2) of the antibacterial activity test of the 15 and 40% aqueous extract of genitri seeds, as well as 50 mg/mL chloramphenicol (positive control), revealed the presence of inhibitory zone formed around the filter paper in the cultures. Inhibition zone diameters of 7.25 and 7.75 mm were observed in the *E. coli* culture when the 15 and 40% of *E. sphaericus* seed extract were respectively tested. For the *L. acidophilus* culture, inhibition zone diameters of 7.5 and 9 mm were recorded for the 15 and 40%, respectively of the *E. sphaericus* seed extract. Meanwhile, there was no inhibition zone observed in the negative control, P1 (distilled water). Table 3 shows the inhibitory effectiveness values of the genitri seed extract on the growth of *E. coli* and *L. acidophilus*. The results indicated that the antimicrobial activity of the seed extract was concentration-dependent. In the *E. coli* culture, a higher value of antibacterial effectiveness was obtained for the 40% concentration (83.78%) compared to the 15% concentration (78.37%). A similar observation was made in the antibacterial culture of *L. Acidophilus*, where values of 87.80% and 73.17% were recorded for the 40 and 15% concentrations, respectively.

The secondary metabolites in genitri seed extract possess inhibitory activities against bacterial growth. Flavonoids are bioactive compounds that have antibacterial properties, and they inhibit cell membrane function by forming a complex with extracellular proteins. They can also damage the bacterial cell membranes to release intracellular compounds.^{34,35} The seeds of genitri contain tannin compounds, which have antibacterial properties.³⁶⁻³⁹ The mechanism of tannin compounds in inhibiting bacterial cells is by denaturing the proteins of the bacterial cells, disrupting the function of the transport system, and synthesis of nucleic acid.⁴⁰ Furthermore, the saponins in the seed extract interact with the bacterial cells and affect the permeability of the cell walls to be ruptured.^{41,42} Moreover, saponins inhibit the activities of enzymes and disrupt bacterial metabolism.⁴³ Alkaloids induced stress in bacteria cells to inhibit their growth.⁴⁴⁻⁴⁶

There are several reports which indicate that the formation of inhibition zones on bacterial growth cultures was induced by flavonoids, glycosides, steroids, alkaloids, saponins, and tannins contents in genitri seeds.⁴⁷⁻⁵⁰ In the present study, the results of the experiments for the two test bacteria showed a significant effect. The diameter and average inhibition zone obtained revealed that the zone of inhibition of *L. acidophilus* was greater than that of *E. coli*. This observation was because *L. acidophilus* is a gram-positive bacteria with a thicker peptidoglycan layer on the cell wall.

Figure 1: Liquid chromatography-mass spectrometry (LC-MS) spectra of *Elaeocarpus sphaericus* seed extractTable 1: Phytochemical constituents of *Elaeocarpus sphaericus* seed extract

Peak	Composition (%)	Mass	Formula	Compound	Class	Peak	Composition (%)	Mass	Formula	Compound	Class
1	0.86	116.0110	C ₄ H ₄ O ₄	Fumaric acid	Dicarboxylic acid	37	1.43	306.0740	C ₁₃ H ₁₄ O ₇	epigallocatechin	Tannin
2	0.78	118.0266	C ₄ H ₆ O ₄	Succinic acid	Dicarboxylic acid	38	2.07	354.0951	C ₁₆ H ₁₈ O ₉	Chlorogenic acid	Tannin
3	0.84	122.0368	C ₇ H ₆ O ₂	Benzoic acid	Aromatic acid	39	1.43	372.1420	C ₁₇ H ₁₂ O ₉	Syringin	Glycoside
4	0.41	116.0837	C ₆ H ₁₂ O ₂	Ethyl butyrate	Ester	40	0.19	402.3498	C ₂₇ H ₄₆ O ₂	δ-tocopherol	Steroid
5	1.27	134.0215	C ₄ H ₆ O ₅	Malic acid	Dicarboxylic acid	41	0.34	416.3654	C ₂₈ H ₄₈ O ₂	γ-tocopherol	Steroid
6	1.45	148.0524	C ₉ H ₈ O ₂	Cinnamic acid	Aromatic acid	42	0.58	400.3705	C ₂₈ H ₄₈ O	Campesterol	Terpenoid
7	1.94	164.0473	C ₉ H ₈ O ₃	p-coumaric acid	Aromatic acid	43	0.85	412.3705	C ₂₉ H ₄₈ O	Isofucosterol	Terpenoid
8	2.16	164.0685	C ₆ H ₁₂ O ₃	Rhamnose	Glucose	44	0.91	412.3705	C ₂₉ H ₄₈ O	24-methylene pollinastanol	Steroid
9	2.27	150.0528	C ₃ H ₁₀ O ₃	Xylose	Glucose	45	0.61	412.3705	C ₂₉ H ₄₈ O	stigmastanol	Steroid
10	1.73	168.0423	C ₈ H ₈ O ₄	Vanillic acid	Aromatic acid	46	0.66	414.3862	C ₂₉ H ₅₀ O	β-sitosterol	Steroid

Peak	Composition (%)	Mass	Formula	Compound	Class	Peak	Composition (%)	Mass	Formula	Compound	Class
11	1.42	178.0266	C ₉ H ₆ O ₄	esculetin	Coumarin	47	1.23	426.3861	C ₃₀ H ₅₀ O	β-amyrin	Terpenoid
12	3.12	180.0423	C ₉ H ₈ O ₄	caffeic acid	Aromatic acid	48	0.75	426.3862	C ₃₀ H ₅₀ O	Cycloeculenol	Terpenoid
13	1.31	192.0423	C ₁₀ H ₈ O ₄	scopoletin	Coumarin	49	0.77	426.3862	C ₃₀ H ₅₀ O	obtusifolol	Terpenoid
14	1.95	194.0579	C ₁₀ H ₁₀ O ₄	Ferulic acid	Carboxylic acid	50	0.82	426.3862	C ₃₀ H ₅₀ O	cycloartenol	Terpenoid
15	0.90	211.1572	C ₁₂ H ₂₁ NO ₂	Elaeocarpine C	Alkaloid	51	0.43	430.3811	C ₂₉ H ₅₀ O ₂	α-tocopherol	Steroid
16	1.59	257.1416	C ₁₆ H ₁₉ NO ₂	(+)-elaecarpine	Alkaloid	52	1.76	432.1058	C ₃₁ H ₂₀ O ₁₀	kaempferol-3-rhamnoside	Flavonoid
17	1.88	257.1416	C ₁₆ H ₁₉ NO ₂	elaecarpine	Alkaloid	53	1.13	440.4018	C ₃₁ H ₅₂ O	24-methylene cycloartanol	Terpenoid
18	1.36	257.1416	C ₁₆ H ₁₉ NO ₂	Isolaecarpine	Alkaloid	54	2.18	448.1006	C ₃₁ H ₂₀ O ₁₁	kaempferol-3-O-D-glucoside	Flavonoid
19	0.86	258.1732	C ₁₆ H ₂₃ N ₂ O	Grandisine B	Alkaloid	55	2.58	448.1006	C ₃₁ H ₂₀ O ₁₁	kaempferol-7-O-β-D-glucoside	Flavonoid
20	1.14	259.1572	C ₁₆ H ₂₁ NO ₂	isolaecarpiline	Alkaloid	56	1.91	448.1006	C ₃₁ H ₂₀ O ₁₁	luteolin-7-glucoside	Flavonoid
21	1.51	261.1729	C ₁₆ H ₂₃ NO ₂	Grandisine D	Alkaloid	57	0.29	450.3498	C ₃₁ H ₄₈ O ₂	phylloquinone	Quinone
22	1.03	267.1735	C ₁₇ H ₂₁ N ₃	elaecarpidine	Alkaloid	58	1.45	484.3189	C ₃₀ H ₄₄ O ₅	Elaecarpucin E	Terpenoid
23	2.26	272.0685	C ₁₃ H ₁₂ O ₅	Naringenin	Flavonoid	59	1.07	484.3189	C ₃₀ H ₄₄ O ₅	Elaecarpucin H	Terpenoid
24	1.89	257.1521	C ₁₆ H ₂₁ NO ₃	isolaecarpicine	Alkaloid	60	1.72	486.3345	C ₃₀ H ₄₆ O ₅	Elaecarpucin A	Terpenoid
25	1.39	276.1838	C ₁₆ H ₂₄ N ₂ O ₂	Grandisine F	Alkaloid	61	1.28	486.3345	C ₃₀ H ₄₆ O ₅	Elaecarpucin B	Terpenoid
26	0.89	277.1678	C ₁₆ H ₂₃ NO ₃	Grandisine A	Alkaloid	62	1.85	486.3345	C ₃₀ H ₄₆ O ₅	Elaecarpucin C	Terpenoid
27	0.98	277.1678	C ₁₆ H ₂₃ NO ₃	Grandisine C	Alkaloid	63	0.94	486.3345	C ₃₀ H ₄₆ O ₅	Elaecarpucin G	Terpenoid
28	1.14	277.1878	C ₁₆ H ₂₃ NO ₃	Grandisine E	Alkaloid	64	0.76	488.3502	C ₃₀ H ₄₈ O ₅	Elaecarpucin F	Terpenoid
29	1.24	279.1834	C ₁₆ H ₂₅ NO ₃	Habbemine A	Alkaloid	65	0.72	516.3087	C ₃₀ H ₄₄ O ₇	Cucurbitacin D	Terpenoid
30	1.46	279.1834	C ₁₆ H ₂₅ NO ₃	Habbemine B	Alkaloid	66	0.84	518.3244	C ₃₀ H ₄₆ O ₇	Cucurbitacin F	Terpenoid
31	2.11	286.0477	C ₁₃ H ₁₀ O ₆	Luteolin	Flavonoid	67	2.33	610.1534	C ₂₇ H ₃₀ O ₁₆	Rutin	Flavonoid
32	2.53	286.0477	C ₁₃ H ₁₀ O ₆	Kaempferol	Flavonoid	68	2.09	624.1690	C ₂₈ H ₃₂ O ₁₆	isorhamnetin-3-O-rutinoside	Flavonoid
33	1.76	290.0790	C ₁₃ H ₁₄ O ₆	Catechin	Tannin	69	1.94	638.1847	C ₂₉ H ₃₄ O ₁₆	rhamnazin-3-rutinoside	Flavonoid
34	0.85	290.1994	C ₁₇ H ₂₆ N ₂ O ₂	Grandisine G	Alkaloid	70	2.53	770.2269	C ₃₄ H ₄₂ O ₂₀	isorhamnetin-3-rutinoside-4'-rhamnoside	Flavonoid
35	1.37	298.0477	C ₁₆ H ₁₀ O ₆	Trifolol	Coumestan	71	1.42	952.0818	C ₄₁ H ₂₆ O ₂₇	Geranin	Tannin
36	2.51	302.0427	C ₁₃ H ₁₀ O ₇	Quercetin	Flavonoid	72	1.94	1110.1033	C ₄₀ H ₃₄ O ₃₂	elaecarpusin	Tannin

Table 2: Zone of inhibition of *Elaeocarpus sphaericus* seed extract against test bacteria

Bacterial Type	Data description	Treatment Type			
		P1	P2	P3	P4
<i>E. coli</i> (F1)	Average diameter of inhibition zone (mm)	0	9.25	7.25	7.75
	Inhibition Zone	0	0.85	0.45	0.55
<i>L. acidophilus</i> (F2)	Average diameter of inhibition zone (mm)	0	10.25	7.50	9.00
	Inhibition Zone	0	1.05	0.50	0.80

P1: Sterile aquadest; P2: 5% chloramphenicol; P3: 15% *Elaeocarpus sphaericus* seed extract; P4: 40% *Elaeocarpus sphaericus* seed extract; F1: *E. coli*; F2: *L. acidophilus*

Table 3: Inhibitory effectiveness of *Elaeocarpus sphaericus* seed extract

Treatment	Effectiveness rate (%)	
	<i>E. coli</i>	<i>L. acidophilus</i>
P3 (15%)	78.37	73.17
P4 (40%)	83.78	87.80

P3: 15% *Elaeocarpus sphaericus* seed extract; P4: 40% *Elaeocarpus sphaericus* seed extract

Even though they have a thick cell wall, gram-positive bacteria generally have a simpler cell wall structure containing 90% peptidoglycan, while the other layer is teichoic acid. This causes the cell walls to be easily damaged by antibacterial compounds. In contrast, *E. coli* is a gram-negative bacterium with a complex cell wall structure that prevent quick denaturing.⁵¹⁻⁵³

Conclusion

The findings of this study indicated that *E. sphaericus* Schum seed extract contains some phytochemical components such as dicarboxylic acid, aromatic acid, ester, glucose, coumarin, alkaloid, flavonoid, tannin, glycoside, steroid, terpenoid, quinone, and coumestan. These compounds have antibacterial activity against *E. coli* and *L. acidophilus*. Therefore, the seed extract of *E. sphaericus* Schum can be potentially used as an antibacterial agent.

Conflict of Interest

The authors declare no conflict of interest.

Authors' Declaration


The authors hereby declare that the work presented in this article is original and that any liability for claims relating to the content of this article will be borne by them.

Acknowledgements

The authors would like to thank the Ministry of Research, Technology and Higher Education of Indonesia for the financial support of this research. Also, the authors are grateful to the Microbiology and Chemistry laboratories for providing the laboratory facilities. Finally, our great appreciation goes to Muhammad Arisandi for helping in data analysis.

References

- Kong JM, Goh NK, Chia LS, Chia TF. Recent advances in traditional plant drugs and orchids. *Acta Pharmacol Sin.* 2003; 24(1):7-21.

**Figure 2:** Antibacterial test of *Elaeocarpus sphaericus* seed extract against *L. acidophilus* (a) and *E. coli* (b)

- Yuan H, Ma Q, Ye L, Piao G. The traditional medicine and modern medicine from natural products. *Molecules.* 2016; 21(5):559.
- Hossen MJ, Uddin MB, Uddin Ahmed SS, Yu ZL, Cho JY. Traditional Medicine/Plants for the Treatment of Reproductive Disorders in Asia Nations. *Pak Vet J.* 2016; 36(2):127-133.
- Pan SY, Litscher G, Gao SH, Zhou SF, Yu ZL, Chen HQ, Zhang SF, Tang MK, Sun JN, Ko KM. Historical perspective of traditional indigenous medical practices: The current renaissance and conservation of herbal resources. *Evidence-based Complement Altern Med.* 2014; 2014(525340):1-20.
- Vasanthi, HR, ShriShriMal N, Das DK. Phytochemicals from Plants to Combat Cardiovascular Disease. *Curr Med Chem.* 2012; 19(14):2242-2251.
- Altemimi A, Lakhssassi N, Baharlouei A, Watson DG, Lightfoot DA. Phytochemicals: Extraction, isolation, and identification of bioactive compounds from plant extracts. *Plants.* 2017; 6(42):1-23.
- Ekor M. The growing use of herbal medicines: Issues relating to adverse reactions and challenges in monitoring safety. *Front Pharmacol.* 2014; 4:1-10.
- Saxena M, Saxena J, Nema R, Singh D, Gupta A. Phytochemistry of medicinal plants. *J Pharmacogn Phytochem.* 2013; 1(6):168-182.
- Panth N, Paudel KR, Karki R. Phytochemical profile and biological activity of *Juglans regia*. *J Integr Med.* 2016; 14(5):359-373.
- Andriani Y, Ramli NM, Syamsunir DF, Kassim MNI, Jaafar J, Aziz NA, Marlina L, Musa NS, Mohamad H. Phytochemical analysis, antioxidant, antibacterial and cytotoxicity properties of keys and cores part of *Pandanus tectorius* fruits. *Arab J Chem.* 2019; 12(8):3555-3564.
- Kanjanasirir P, Suksatu A, Manopwisedjaroen S, Munyoo

- B, Tuchinda P, Jearawuttanakul K, et al. High-content screening of Thai medicinal plants reveals *Boesenbergia rotunda* extract and its component Panduratin A as anti-SARS-CoV-2 agents. *Sci Rep*. 2020; 10(1):1-12.
12. Ayaz M, Ullah F, Sadiq A, Ullah F, Ovais M, Ahmed J, Devkota HP. Synergistic interactions of phytochemicals with antimicrobial agents: Potential strategy to counteract drug resistance. *Chem Biol Interact*. 2019; 308(May):294-303.
 13. Omojate GC, Enwa FO, Jewo AO, Eze CO. Mechanisms of Antimicrobial Actions of Phytochemicals against Enteric Pathogens. *J Pharm Chem Biol Sci*. 2014; 2(2):77-85.
 14. Azih I, Ijezie M, Ugariogu SN, Akalezi CO. Antibacterial activity and identification of metabolites from the semi-purified fraction of *Chrysophyllum albidum* leaf (African star apple). *Trop J Nat Prod Res*. 2020; 4(7):262-269.
 15. Rohandi A, Gunawan. Distribution and the potential growth of ganitri (*Elaeocarpus ganitrus* Roxb) in Central Java. *Jurnal Ilmu Kehutanan*. 2014; 8(1):25-33.
 16. Nain J, Garg K, Dhahiya S. Analgesic and anti-inflammatory activity of *Elaeocarpus sphaericus* leaf extract. *Int J Pharm Pharm Sci*. 2012; 4(SUPPL.1):379-381.
 17. Hule AK, Shah AS, Gambhire MN, Juvekar AR. An evaluation of the antidiabetic effects of *Elaeocarpus ganitrus* in experimental animals. *Indian J Pharmacol*. 2011; 43(1):56-59.
 18. Joshi S, Gupta P, Kumar N, Rai N, Gautam P, Thapliyal A. A comprehensive report on therapeutic potential of *Elaeocarpus ganitrus* Roxb.(Rudraksha). *Environ Conserv J*. 2012; 13(3):147-150.
 19. Ogundele AV, Das AM. Chemical Constituents from the Leaves of *Elaeocarpus floribundus* Ayorinde. *Nat Prod Res*. 2021; 35(3):517-520.
 20. Tripathy S, Mida A, Swain SR. Phytochemical Screening and Thin Layer Chromatographic Studies of *Elaeocarpus ganitrus* Seed the Magical Electromagnetic Bead (Rudraksha). *Int J Pharm Biol Sci*. 2016; 6(3):16-24.
 21. Dalei J, Sahoo D. Evaluation of antimicrobial activity and phytochemical screening of epicarp and endocarp parts of *Elaeocarpus ganitrus*. *Int J Pharm Bio Sci*. 2016; 7(2):265-9.
 22. Jain PK, Sharma P, Joshi SC. Antioxidant and lipid lowering effects of elaeocarpus ganitrus in cholesterol fed rabbits. *Int J Pharmaceut Sci and Res*. 2018; 9(2):526-534.
 23. Bhatt BD, Dahal P. Antioxidant and Antimicrobial Efficacy of Various Solvent Extracts of Seed of Rudraksha (*Elaeocarpus ganitrus*) from Ilam District of Nepal. *J Nepal Chem Soc*. 2019; 40:11-18.
 24. Poolman JT, Wacker M. Extraintestinal pathogenic *Escherichia coli*, a common human pathogen: challenges for vaccine development and progress in the field. *J Infect Dis*. 2016; 213(1):6-13.
 25. Lin CK, Tsai HC, Lin PP, Tsen HY, Tsai CC. *Lactobacillus acidophilus* LAP5 able to inhibit the *Salmonella choleraesuis* invasion to the human Caco-2 epithelial cell. *Anaerobe*. 2008; 14(5):251-255.
 26. Wu Z, Yang K, Zhang A, Chang W, Zheng A, Chen Z, Cai H, Liu G. Effects of *Lactobacillus acidophilus* on the growth performance, immune response, and intestinal barrier function of broiler chickens challenged with *Escherichia coli* O157. *Poult Sci*. 2021; 100(9):101323.
 27. Balouiri M, Sadiki M, Ibensouda SK. Methods for *in vitro* evaluating antimicrobial activity: A review. *J Pharm Anal*. 2016; 6(2):71-79.
 28. Souiri D, Salimi N, Ghabooli M. Hydrothermal fabrication of pure ZnSe nanocrystals at different microwave irradiation times and their disc-diffusion antibacterial potential against Gram negative bacteria: Bio-optical advantages. *Inorg Chem Commun*. 2021; 123:108345.
 29. Dwivedi P, Narvi SS, Tewari RP. Phytofabrication characterization and comparative analysis of Ag nanoparticles by diverse biochemicals from *Elaeocarpus ganitrus* Roxb., *Terminalia arjuna* Roxb., *Pseudotsuga menziesii*, *Prosopis spicigera*, *Ficus religiosa*, *Ocimum sanctum*, *Curcuma longa*. *Ind Crops Prod*. 2014; 54:22-31.
 30. Wilapangga A and Syaputra S. Antibacterial analysis of agar disc method and toxicity test using BSLT (Brine Shrimp Lethality Test) from methanol extract of Bay leaf (*Eugenia polyantha*). *Indones J Biotechnol Biodivers*. 2018;2(2):50-56.
 31. Nurfitriani R, Krishanti N putu RA, Akhdiya A, Wahyudi AT. Filospheric bacteria screening produce anti *Xanthomonas oryzae* bioactive compounds pv. oryzae causes bacterial leaf blight in rice. *J Sumberd Hayati*. 2016; 2(1):19-24.
 32. Hamzah A. *In Vitro* Analysis of Dragon Scale Leaves (*Drymoglossum pilosellaoides*) Antibacterial Activity Against *Vibrio harveyi* and *Vibrio parahaemolyticus* Bacteria. *J Aquac Fish Heal*. 2019; 8(2):86-91.
 33. Pouvoa D, Kokoskova B, Pavela R, Rysanek P. Effectivity of plant essential oils against *Clavibacter michiganensis*, *in vitro*. *Zemdirbyste*. 2008; 95(3):440-446.
 34. Salem MZM, Ali HM, El-Shanhorey NA, Abdel-Megeed A. Evaluation of extracts and essential oil from *Callistemon viminalis* leaves: Antibacterial and antioxidant activities, total phenolic and flavonoid contents. *Asian Pac J Trop Med*. 2013; 6(10):785-791.
 35. Tagousop CN, Tamokou J-D, Ekom SE, Ngokam D, Voutquenne-Nazabadioko L. Antimicrobial activities of flavonoid glycosides from *Graptophyllum grandulosum* and their mechanism of antibacterial action. *BMC Complement Altern Med*. 2018; 18(1):1-10.
 36. Banso A and Adeyemo SO. Evaluation of antibacterial properties of tannins isolated from *Dichrostachys cinerea*. *African J Biotechnol*. 2007; 6(15):1785-1787.
 37. Doss A, Mubarak HM, Dhanabalan R. Antibacterial activity of tannins from the leaves of *Solanum trilobatum* Linn. *Indian J Sci Technol*. 2009; 2(2):41-43.
 38. Mailoa MN, Mahendradatta M, Laga A, Djide N. Antimicrobial Activities Of Tannins Extract From Guava Leaves (*Psidium Guajava* L) On Pathogens Microbial. *Int J Sci Technol Res*. 2014; 3(1):236-241.
 39. Liu M, Yang K, Wang J, Zhang J, Qi Y, Wei X, Fan M. Young astringent persimmon tannin inhibits methicillin-resistant *Staphylococcus aureus* isolated from pork. *LWT*. 2019; 100(July 2018):48-55.
 40. Javed B, Nawaz K, Munazir M. Phytochemical Analysis and Antibacterial Activity of Tannins Extracted from *Salix alba* L. Against Different Gram-Positive and Gram-Negative Bacterial Strains. *Iran J Sci Technol Trans A Sci*. 2020; 44(5):1303-1314.
 41. Mandal P, Babu SPS, Mandal NC. Antimicrobial activity of saponins from *Acacia auriculiformis*. *Fitoterapia*. 2005; 76(5):462-465.
 42. Khan MI, Ahmed A, Shin JH, Baek JS, Kim MY, Kim JD. Green Tea Seed Isolated Saponins Exerts Antibacterial Effects against Various Strains of Gram Positive and Gram Negative Bacteria, a Comprehensive Study *in vitro* and *in vivo*. *Evidence-based Complement Altern Med*. 2018; 2018(3486106):1-12.
 43. Mabhiza D, Chitemerere T, Mukanganyama S. Antibacterial Properties of Alkaloid Extracts from *Callistemon citrinus* and *Vernonia adoensis* against *Staphylococcus aureus* and *Pseudomonas aeruginosa*. *Int J Med Chem*. 2016; 2016(6304163):1-7.
 44. Avcı FG, Sayar NA, Sariyar Akbulut B. An OMIC approach to elaborate the antibacterial mechanisms of different alkaloids. *Phytochemistry*. 2018; 149:123-131.
 45. Jain A, Parihar DK. Antibacterial, biofilm dispersal and antibiofilm potential of alkaloids and flavonoids of Curcuma. *Biocatal Agric Biotechnol*. 2018; 16:677-682.
 46. Zhou LN, Ge XL, Dong TT, Gao HY, Sun BH.

- Antibacterial steroidal alkaloids from *Holarrhena antidysenterica*. Chin J Nat Med. 2017; 15(7):540-545.
47. Hernández NE, Tereschuk ML, Abdala LR. Antimicrobial activity of flavonoids in medicinal plants from Tafi del Valle (Tucuman, Argentina). J Ethnopharmacol. 2000; 73(1-2):317-322.
48. Mamtha B, Kavitha K, Srinivasan KK, Shivananda PG. An in vitro study of the effect of *Centella asiatica* [Indian pennywort] on enteric pathogens. Indian J Pharmacol. 2004; 36(1):41-42.
49. Mujeeb F, Bajpai P, Pathak N. Phytochemical evaluation, antimicrobial activity, and determination of bioactive components from leaves of aegle marmelos. Biomed Res Int. 2014; 2014(497606):1-11.
50. Ali M, Yahaya A, Zage A, Yusuf Z. In vitro Antibacterial Activity and Phytochemical Screening of Psidium guajava on Some Enteric Bacterial Isolates of Public Health Importance. J Adv Med Pharm Sci. 2017; 12(3):1-7.
51. Kline KA, Lewis AL. Gram-Positive Uropathogens, Polymicrobial Urinary Tract Infection, and the Emerging Microbiota of the Urinary Tract. Microbiol Spectr. 2016; 4(2):139-148.
52. Urmann K, Arshavsky-Graham S, Walter JG, Scheper T, Segal E. Whole-cell detection of live: *Lactobacillus acidophilus* on aptamer-decorated porous silicon biosensors. Analyst. 2016; 141(18):5432-5440.
53. Lingga AR, Pato U, Rossi E. Antibacterial test of kecombrang (*Nicolaia speciosa* horan) stem extract against *Staphylococcus aureus* and *Escherichia coli*. JOM Faperta. 2016; 3(1):1-15.

ORIGINALITY REPORT

10%

SIMILARITY INDEX

5%

INTERNET SOURCES

11%

PUBLICATIONS

5%

STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

3%

★ R C Megananda, Y N Azhizhah, Pujiati, C N Primiani. "Oil content analysis on yam bean fermented by *Aspergillus niger*", Journal of Physics: Conference Series, 2020

Publication

Exclude quotes On

Exclude bibliography On

Exclude matches < 20 words