

Characteristics and Seasonal Abundance of Seagrass at Pidakan Coast Pacitan, East Java, Indonesia

by Nurul Kusuma Dewi, Joko Widiyanto

Submission date: 06-Feb-2019 11:50PM (UTC-0800)

Submission ID: 1074385927

File name: 14._II.A.c.1.a.1_PROSIDING_HCU.pdf (1.97M)

Word count: 4001

Character count: 22741

Characteristics and Seasonal Abundance of Seagrass at Pidakan Coast Pacitan, East Java, Indonesia

23

Nurul Kusuma Dewi^{1*} and Joko Widiyanto²

^{1,2}Biology Education Study Program, Faculty of Teacher Training and Education, Universitas PGRI Madiun

*Corresponding E-mail: nurulkd@unipma.ac.id

Abstract

This research was conducted to study the condition of seagrass at Pidakan Coast Pacitan, East Java, Indonesia during rainy and dry season using a percentage of the covered area and physico chemical factors. The method used was transect-plot. The physical and chemical factors measured included: light penetration, high water level, current velocity, salinity, temperature, wave height, DO, pH, sediment texture, and nutrients of NH₄, NO₃, P₂O₅, C-organic. The result showed that seagrass bed in Pidakan Coast was formed by only single species, *Thalassia hemprichii*. The average percentage of seagrass cover was 30.89% in the rainy season and 15.05% in the dry season. According to decree of the Minister of Environment No. 200 in 2004, the seagrass bed conditions in Pidakan Coast in the rainy season was categorized into less rich/less healthy, whereas in the dry season was categorized into poor. The factors that influence the distribution and abundance of seagrass in the Pidakan Coast were substrate, depth, and the waves. The decline in the seagrass covering percentage in the dry season was caused by light and temperature factors. High light and temperatures caused dieback of seagrass. Seagrasses in Pidakan Coast were associated with different types of organisms.

Keywords: Dieback; Light; Seagrass; Season; Temperature

6

1. Introduction

Seagrass, the marine flowering plants capable of completing its lifecycle fully submerged underwater, represents a dominant biological community along the shallow coastal areas of the world. Seagrass is one of the most important components of tropical coastal ecosystems, besides mangroves and coral reefs (de la Torre-Castro, 2006), and has wide global distribution (Hartog, 1970). Compared with mangroves and coral reefs, the seagrass ecosystem has perhaps received less scholarly attention, yet this ecosystem has much potential benefit and provides crucial services (de la Torre Castro et al., 2014).

Seagrasses provide a wide range of ecosystem services (Campagner et al., 2015; Cullen-Unsworth et al., 2014; Nordlund et al., 2016) here defined as natural processes and components that directly or indirectly benefit human needs (de Groot et al., 2002). Seagrass meadows are highly productive ecosystems that offer several ecosystem goods and services such as food, recreation, natural shelter, conservation, among others (Tuya et al., 2014). This habitat is used for shelter, foraging, spawning, and rearing place for marine organisms (Hemminga and Duarte, 2003). Seagrass beds are also positioned very close to the mainland, thereby, is influenced by anthropogenic effect (Ambo-Rappe, 2014).

Seagrasses are generally assigned to two families, Potamogetonaceae and Hydrocharitaceae encompassing 12 genera of angiosperms containing about 50 species. Seagrasses occur in all coastal areas of the world, except along Antarctic shores. Seagrass beds are common in shallow waters of the tropics, subtropics, and temperate regions (Green and Short, 2003). Seagrass species are often separated into tropical and temperate genera. The former are considered to comprise seven of the genera, while the latter comprise the remaining five genera (Hemminga and Duarte, 2003).

Most seagrass meadows are monospecific, particularly those in the temperate zone, but this holds also for tropical and subtropical areas with multispecific floras. Although seagrass

species richness, the number of species present in any one meadow, can be significant, seagrass species diversity, i.e. the evenness of their contribution to the community, is typically low, as even in tropical multispecific meadows the distribution of biomass is generally skewed, with one or a few species comprising most of the biomass of the community (Terrados and Ros, 1991). The major part of the biological diversity of seagrass meadows, however, is contributed by the rich associated fauna and algal floras, and not by the seagrasses themselves. The meadows with the richest species diversity are found in the Indo-Pacific area and the Red Sea, where mixed meadows are abundant, containing up to 12 co-occurring species (Duarte, 2000).

The distribution and stability of seagrass community is determined by factors such as: nutrients, light, sediment, salinity, and temperature (Udy and Dennison 1997, Ralph et al., 2007; Hemminga and Duarte 2000; Kahn and Dur, 2006; Masini et al., 1995; Campbell et al., 2006). The influential nutrients are nitrogen (N) and phosphorus (P). The growth of *H. uninervis* is limited by N, *Zostera capricorni* Aschers is limited by N and P; otherwise *C. serrulata* are not affected by the addition of any nutrients (Udy and Dennison, 1997).

The enrichment of nutrients (eutrophication) can trigger the growth of epiphytic algae on the leaves of seagrass and algae in the water column. Both types of blooming algae reduce the amount of light reaching the seagrass. This reduces the effectiveness of photosynthesis so as to reduce the productivity of seagrasses and leads to the decline of seagrass communities worldwide (Hays, 2005; Waycott et al., 2009). Certain light intensity is important for seagrasses for photosynthesis (Ralph et al., 2007). Masini et al. (1995) reported that the distribution of seagrass species is related to the need for light. Reduced light penetration into seagrass waters other than caused by eutrophication can also be caused by increased sediment load (Ralph et al. 2007).

Seagrasses can grow in various habitat types, but most species of seagrass grow on sand and silt substrate. This is because the habitat is more easily penetrated by the roots of seagrass. The seagrass species have different salinity tolerances, but most have a wide range of tolerances to salinity between 10-440/00 (Hemminga and Duarte, 2000). Low or high temperature pressures cause disturbances to photosynthesis and the decline of seagrass biomass (Masini et al., 1995; Campbell et al., 2006). Some species of tropical seagrass, *C. rotundata*, *C. serrulata*, *H. uninervis* and *T. hemprichii* are more tolerant of high temperature pressures than *H. ovalis*, *Z. capricorni* and *S. isoetifolium* (Campbell et al., 2006).

In the Philippines, daytime temperature and length are the most influential factors in density, biomass and seagrass production. Maximum daylight poses a negative effect on seagrass. While Erftemeijer and Herman (1994) report, in the tropical region of Gusung Tallang, South Sulawesi, the biomass, production, and nutrient content of the *E. acoroides* seawater network do not respond to rainfall, salinity, tidal, nutrient availability, and water movement or turbidity. This shows that the watershed forest in the area is still healthy so it can be assumed that the nutrient input is the same between the dry and rainy seasons.

Indonesia has two seasons throughout the year, namely dry season (dry) and rain (wet). The two seasons did not take place simultaneously in the Indonesian archipelago. Generally, the dry season occurs in April continuing until about September-October, and the wet season occurring in October-November continues March-May. The islands of Java and Nusa Tenggara experienced a dry season in April-October.

Indonesia is a home to exceptional seagrass diversity and there is evidence that seagrass there is experiencing some decline due to a quick economical development with unavoidable increase of pollution (Kuriandewa et al., 2003), although relatively little is known about the ecology of seagrass compared to other regions of the world (Waycott et al., 2009; Short et al., 2011). Tropical seagrasses, however, can show large temporal changes. For example, the

17 mass and growth of intertidal and upper subtidal seagrass meadows in Southeast (SE) Asia can vary by up to a factor of four during the year (Brouns, 1985; Estacion and Fortes, 1988; Erftemeijer and Herman, 1994; Lanyon and Marsh, 1995; Ethirmannasingam et al., 1996). Tidal exposure and water motion are considered the main environmental factors driving the temporal variation of biomass and growth of tropical seagrasses (Erftemeijer and Herman, 1994), which may lead to more complex temporal patterns than those encountered in temperate seagrasses, where light and temperature are the main environmental factors driving temporal change (Terrados and Ros, 1991; Perez and Romero, 1992). Changes in sea level, salinity, temperature, atmospheric CO₂, and UV radiation can alter seagrass distribution, productivity, and community composition. In turn, potential changes in distribution and structure of seagrass communities may have profound implications for local and regional biota, nearshore geomorphology, and biogeochemical cycles. Few investigations to date have focused specifically on potential responses of seagrasses to the changing global environment (Short and Neckles, 1999). Our goals were to describe the temporal changes in the abundance of *Thalassia hemprichii* (Ehrenb.) Aschers in a shallow intertidal Indonesia seagrass meadow. We characterized the seagrass in the Pidakan Coast, which is located in Pacitan, East Java. At Pidakan Coast, seagrass growing on the dead coral habitat. Information about seagrass in Pacitan is limited. The objective of this research was to study on the condition of seagrass at Pidakan Coast Pacitan, East Java, Indonesia during rainy and dry season using a percentage of the covered area and physico chemical factors.

2. Materials and methods

The method used was transect-plot. We used 5 transect (each transect 50 m) perpendicular to the shoreline. The distance between transect 25 m. At each transect, data was collected using the quadrat plot, 0.5 m x 0.5 m, the distance between plot 5 m. The total number of plots is 55. On each plot, we identified the species, noted percentage of the covered area, and noted other organisms found. The determination of a percentage of the covered area follows percent cover standards from seagrasswatch.org. The physical and chemical factors measured included: light penetration, high water level, current velocity, salinity, temperature, wave height, DO, pH, sediment texture, and nutrients of NH₄, NO₃, P₂O₅, C-organic. Identification of seagrass based on Waycott et al. (2009). There is no statistical analysis was used in this study.

3. Results

Seagrass meadow in Pidakan Coast was formed by only single species, *Thalassia hemprichii*. The average percentage of seagrass cover as below:

Table 1. The average percentage of seagrass cover

Transect	C (percentage of the covered area) in rainy season	C (percentage of the covered area) in dry season
I	22,14 %	14,63 %
III	39,14 %	7,72 %
IV	15 %	3,45 %
V	25,33 %	20,45 %
Average	30,89 %	15,05 %

From the table above, we can see that the average percentage of seagrass cover varies between transects. This is because dispersion and condition of physico-chemical factors also

differ between transects. The average percentage of seagrass cover in the rainy season is higher than in the dry season.

Physico-chemical factors at Pidakan Coast as follows:

Table 2. Physico-chemical factors at Pidakan Coast

No.	Parameter	Satuan	Rainy	Dry
1	Light penetration	m	2,5	3
2	Air temperature	°C	21	31
3	Water temperature	°C	22	33
4	Salinity	‰	39.56	40.05
5	DO		6.4	7.5
6	pH	-	7.795	8.210
7	Current velocity	m/s	0,125	0.1
8	NH ₄	me/l	0.075	0.125
9	NO ₃	me/l	0.04285	0.03345
10	PO ₄	ppm	0.135	0.250
11	Organic C	%	0.40	0.55
12	Organic matter	%	0.69	0.75

The table above shows that physico-chemical factors in the rainy and dry seasons show no significant difference. All the chemical physics and nutrient factors measured are still within the tolerance range of the seagrass.

4. Discussion

Seagrass meadow in Pidakan Coast was formed by only single species, *Thalassia hemprichii*, a dominant meadow-forming species on sediments associated with coral reefs. Although *Thalassia hemprichii* dominates more reefal habitats, it is found on a diversity of substrata, although is limited in areas with any freshwater runoff. *Thalassia hemprichii* (turtlegrass) occurs throughout the Indo-West Pacific although not as far east as the Hawaiian Islands and Fiji. *Thalassia hemprichii* is an important food resource for green turtles in the tropics. This species often dominates the landward edge of islands and cays where it plays a major role in stabilizing sediments with its intertwining rhizome mat (Waycott et al., 2009)

Thalassia hemprichii found from the shallow subtidal to deeper than 10 m, this species does not tolerate long periods of exposure. But, in many places of Indonesia includes Pidakan coast, *T. hemprichii* a seagrass species that is most resistant to environmental stress conditions. *Thalassia hemprichii* also forms dense meadows and has a high productivity. Numerous grazers and other fauna inhabit these meadows. Macroinvertebrates found are 15 species namely *Nerita chamaeleon*, *Trochus niloticus*, *Thais aculeate*, *Cypraea eglantina*, *Portunus pelagicus*, *Tripneustes gratilla*, *Echinometra mathaei*, *Diadema setosum*, *Holothuria atra*, *Ophiomastix xannulosa*, *Ophiomastix variabilis*, *Archaster typicus*, *Ophiarachna affinis*, *Turbo argyrostomus* and *Tetracrita* sp. Most of the macroinvertebrate come from Mollusca Phylum. There are also 16 species of macroalgae found on Pidakan Coast namely: *Ulva lactuca*, *Enteromorpha intestinalis*, *Valoni aegagropila*, *Padina australis*, *Dictyopteris undulata*, *Sargassum binderi*, *Hormophysa triquetra*, *Turbinaria ornata*, *Euclima edule*, *Euclima cottonii*, *Euclima spinosum*, *Amphiroa rigida*, *Jania*

rubens, *Galaxaura rugosa*, *Gelidium amansii* and *Actinotrichia fragilis*. The species found are dominated by Rhodophyta Division.

Thalassia hemprichii is the Indo-West Pacific representative of a genus with only two species, the other, *Thalassia testudinum* occurs in the Caribbean. *Thalassia hemprichii* is recognized by having curved leaves often with tannin cells which look red, purple or dark brown. These cells give the leaf a slightly speckled appearance. This species also has characteristic rhizomes, thick (usually white or pale pink) with triangular shaped scars, and the membranous fibres at each node. *Thalassia hemprichii* has key characteristics as follows: strap-like leaves arise from an upright stem with a fully enclosing leaf sheath. Leaf tip blunt with minute serrations. Rhizome thick with obvious node scars usually triangular in shape and with persistent leaf sheaths. Found throughout the tropical Indo-West Pacific. About reproduction, *Thalassia hemprichii* has separate male and female plants. The flowers form in the regions at the base of the shoot and are obscured by the sheath until they emerge. The male flower elongates on a long stalk (pedicel), the flower when mature separates into six or more parts. The female flower appears similar although has a finer texture, the ovary being at the base of the shoot. Fruits have an almost prickly appearance and may contain up to nine seeds (around 0.6 mm diameter) (Waycott et al., 2009). The appearance of *Thalassia hemprichii* leaves differs between the rainy season and the dry season. In the rainy season, these leaves look fresh green, while in the dry season the leaves of *Thalassia hemprichii* appear burnt and blackened. This is the effect of high radiation of sunlight and temperature.

According to decree of the Minister of Environment No. 200 in 2004, the seagrass bed conditions in Pidakan Coast in the rainy season was categorized 12 to less rich/less healthy, whereas in the dry season was categorized into poor. The factors that influence the distribution and abundance of seagrass in the Pidakan Coast were substrate, depth, and the waves. Substrate in Pidakan Coast is dominated by dead coral, only certain species and sturdy stems that can grow in these habitats. Sediment composition or substrate can also affect the availability of phosphate. In the carbonate sediments (sedimentary minerals derived from coral reefs), phosphate interact with carbonates and less available as a free phosphate. But it is still helped by excess seagrass that can take nutrients such as ammonia and phosphate through the leaves.

Depth is also one factor that influences the distribution and abundance of seagrass in the Pidakan Coast. In Pidakan Coast, seagrass grows only in shallow areas such as reef (dead coral debris). Distribution of seagrass on the beach is limited to a distance of ± 50 m from the shoreline. After that distance, depth immediately increased dramatically. This is why seagrasses are no longer found because sunlight cannot reach up to the base thus blocking seagrass make the process of photosynthesis.

The waves at Pidakan Coast are strong. Strong tidal currents cause difficult seagrass taken root at the bottom of the water so that less breed well. Strong waves also make seagrass vegetation can be ripped from the substrate. Seagrass leaves clean of algae epiphytic showed relatively strong local currents.

The decline in the seagrass covering percentage in the dry season was caused by light and temperature factors. High light and temperature caused dieback of seagrass. Thermal optima associated with high rates of photosynthesis and growth ranges from approximately 15 to 33°C, with the species-specific optimums generally reflecting their geographic distribution (Berry and Bjorkman, 1980; Bulthuis, 1987; Coles and Jokiel, 1977; Collier et al., 2011; Masini and Manning, 1997; Perez and Romero, 1992). During low tide, these thermal optima are likely exceeded for shallow water habitats. Extreme events occurring during low tide have been linked to seagrass loss (Massa et al., 2009; Rasheed and Unsworth, 2011). Although coastal water temperatures are still within the range of seagrass tolerance, but seagrass has

been shown to change their characteristic (morphological appearance) and percent of the covered area.

Seagrasses are affected by thermal stress in a number of ways. Water temperature affects the balance between carbon uptake (photosynthesis) and carbon consumption (respiration) (Bulthuis, 1987; Perez and Romero, 1992). The photosynthetic apparatus is highly sensitive to temperature, with temperature sensitive processes occurring throughout the photosynthetic pathways (Buxton et al., 2012; Jones et al., 1998). Increases in photosynthesis occur within their physiological optimum, but this is followed by sharp reductions in photosynthetic efficiency after temperatures exceed optimum thresholds (Bulthuis, 1983; Campbell et al., 2006; Perez and Romero, 1992; Ralph et al., 1998). Sensitivity to elevated temperature is variable among species, with tropical species tolerating higher temperatures than subtropical and temperate species (Campbell et al., 2006; Collier et al., 2011). Following extreme or prolonged thermal stress, seagrass mortality results; however, *in situ* observations of seagrass mortality typically occur after the thermal event, making it difficult to directly link changes in physiological processes to mortality (Marbá and Duarte, 2010; Massa et al., 2009; Rasheed and Unsworth, 2011).

5. Conclusion

Seagrass meadow in Pidakan Coast was formed by only single species, *Thalassia hemprichii*, a seagrass species that is most resistant to environmental stress conditions. The appearance of *Thalassia hemprichii* leaves differs between the rainy season and the dry season. In the rainy season, these leaves look fresh green, while in the dry season the leaves of *Thalassia hemprichii* appear burnt and blackened. This is the effect of high radiation of sunlight and temperature. The average percentage of seagrass cover was 30.89% in the rainy season and 15.05% in the dry season. The decline in the seagrass covering percentage in the dry season was caused by light and temperature factors. High light and temperatures caused dieback of seagrass. Seagrasses in Pidakan Coast were associated with different types of organisms.

References

- Ambo-Rappe, R. (2014). Developing a methodology of bioindication of human-induced effected using seagrass morphological variation in Spremonde Archipelago, South Sulawesi, Indonesia. *Marine Pollution Bulletin*, 86, 298-303.
- Berry, J., & Bjorkman, O. (1980). Photosynthetic response and adaptation to temperature in higher plants. *Annual Review of Plant Physiology Journal*, 31, 491-543.
- Brouns, J. J. (1985). A Comparison of the annual production and biomass in three monospecific stands of the seagrass *Thalassia hemprichii* (Ehrenb.) aschers. *Aquatic Botany*, 23, 149-175.
- Bulthuis, D. (1983). Effects of temperature on the photosynthesis-irradiance curve of the Australian seagrass, *Heterozostera tasmanica*. *Marine biology letters*. 4, 47-57.
- Bulthuis, D. A. (1987). Effect of temperature on photosynthesis and growth of seagrasses. *Aquatic Botany*, 27, 27-40.
- Buxton, L., Takahashi, S., Hill, R., & Ralph, P. (2012). Variability in the primary site of photosynthetic damage in *Symbiodinium* sp. (Dinophyceae) exposed to thermal stress. *Journal of Phycology*, 48, 117-126.
- Campagne, C. S., Salles, J. M., Boissery, P., & Deter, J. (2015). The seagrass *Posidonia oceanica*: ecosystem services identification and economic evaluation of goods and benefits. *Marine Pollution Bulletin*, 97, 391-400.
- Campbell, S., McKenzie, L., & Kerville, S. (2006). Photosynthetic responses of seven tropical seagrasses to elevated seawater temperature. *Journal of Experimental Marine Biology and Ecology*, 330, 455-468.

- Coles, S., & Jokiel, P. (1977). Effects of temperature on photosynthesis and respiration in hermatypic corals. *Marine Biology*, 43, 209-216.
- Collier, C., Uthicke, S., & Waycott, M. (2011). Thermal tolerance of two seagrass species at contrasting light levels: implications for future distribution in the great barrier reef. *Limnology and Oceanography*, 56, 2200-2210.
- Cullen-Unsworth, L., Mtwana, N., Paddock, J., Baker, S., McKenzie, L., & Unsworth, R. (2014). Seagrass meadows globally as a coupled social-ecological system: implications for human well being. *Marine Pollution Bulletin*, 83, 387-397.
- de Groot, R. S., Wilson, M. A., & Boumans, R. M. (2002). A Typology for the classification, description, and valuation of ecosystem functions, goods and services. *Ecological Economics*, 41, 393-408.
- de la Torre Castro, M., Di Carlo, G., & Jiddawi, N. (2014). Seagrass Importance for a small scale fishery in the tropics: the need for seascape management. *Marine Pollution Bulletin*, 83, 398-347.
- de la Torre-Castro, M. (2006). *Humans and seagrass in east Africa - a socialecological system approach*. Stockholm: Department of Systems Ecology, Stockholm University.
- Duarte, C. M. (2000). Marine biodiversity and ecosystem services an elusive link. *Journal of Experimental Marine Biology and Ecology*, 250, 117-131.
- Ertfemeijer, P. L., & Herman, P. M. (1994). Seasonal changes in environmental variables, biomass, production, and nutrient contents in two contrasting tropical intertidal seagrass beds in south Sulawesi (Indonesia). *Oecologia*, 99, 45-59.
- Estacion, J. S., & Fortes, M. D. (1988). Growth rates and primary production of *Enhalus acoroides* (L.f.) royle and its epiphytes. *Aquatic Botany*, 29, 347-356.
- Ethirnanasingam, S., Phang, S. M., & Sasekumar, A. (1996). A study of some phenological events in a Malaysian *Enhalus acoroides* bed. In: Kuo, J., Philips, R.C., Walker, D., Kirkman, H. (Eds.). *Seagrass biology proceedings of an international workshop* (pp. 33 - 40). Rottneest Island, Western Australia: The Faculty of Science, University of Western Australia.
- Green, E., & Short, F. (2003). *World atlas of seagrass*. Berkeley, USA: University of California Press.
- Hartog, D. (1970). *The Sea-grasses of the world*. Amsterdam, North Holland: Verhandeligen der Koninklijke Nederlandse Akademie van Wetenschappen, afd. Natuurkunde, Reek 2.
- Hemminga, M. A., & Duarte, C. M. (2003). *Seagrass Ecology*. Cambridge: Cambridge University Press.
- Jones, R., Hoegh-Guldberg, O., Larkum, A., & Schreiber, U. (1998). Temperature-induced bleaching of corals begins with impairment of the CO₂ fixation mechanism in zooxanthellae. *Plant Cell Environ*, 21, 1219-1230.
- Kuriandewa, T. E., Kiswara, E., Hutomo, M., & Soemodihardjo, S. (2003). The Seagrasses of Indonesia. . In E. S. Green, *World Atlas of Seagrass* (pp. 171 - 182). Berkeley, USA: University of California Press.
- Lanyon, J. M., & Marsh, H. (1995). Temporal changes in the abundance of some tropical intertidal seagrasses in North Queensland. *Aquatic Botany*, 49, 217-237.
- Marbá, N., & Duarte, C. (2010). Mediterranean warming triggers seagrass (*Posidonia oceanica*) shoot mortality. *Global Change Biology*, 16, 2366-2375.
- Masini, R., & Manning, C. (1997). The photosynthetic responses to irradiance and temperature of four meadow-forming seagrasses. *Aquatic Botany*, 58, 21-36.
- Massa, S., Arnaud-Haond, S., Pearson, G., & Serrão, E. (2009). Temperature tolerance and survival of intertidal populations of the seagrass *Zostera noltii* (Hornemann) in southern Europe (Ria Formosa, Portugal). *Hydrobiologia*, 619, 195-201.

- Nordlund, L. M., Koch, E. W., Barbier, E. B., & Creed, J. C. (2016). Seagrass ecosystem services and their variability across genera and geographical regions. *PloS One*, 2016.
- Perez, M., & Romero. (1992). Photosynthetic response to light and temperature of the seagrass *Cymodocea nodosa* and the prediction of its seasonality. *Aquatic Botany*, 43, 51-62.
- Ralph, P., Gademann, R., & Dennison, W. (1998). *In situ* seagrass photosynthesis measured using a submersible, pulse-amplitude modulated fluorometer. *Marine Biology*, 132, 367-373.
- Rasheed, M., & Unsworth, R. (2011). Long-term climate-associated dynamics of a tropical seagrass meadow: implications for the future. *Marine Ecology Progress Series*, 422, 93-103.
- Short, F. T., & Neckles, H. A. (1999). The effects of global climate change on seagrasses. *Aquatic Botany*, 63, 169-196.
- Short, F. T., Polidoro, B., Livingstone, S. R., & Carpenter, K. E. (2011). Extinction risk assessment of the world's seagrass species. *Biological Conservation*, 144, 1961-1971.
- Terrados, J., & Ros, J. (1991). Production dynamics in a macrophyte-dominated ecosystem. *Oecologia aquatica*, 10, 255-270.
- Tuya, F., Haroun, R., & Espino, F. (2014). Economic assessment of ecosystem services: monetary value of seagrass meadows for coastal fisheries. *Ocean and Coastal Management*, 96, 181-187.
- Waycott, M., Duarte, C. M., Carruthers, T. J., & Orth, R. J. (2009). Accelerating loss of seagrass across the globe threatens coastal ecosystems. *PNAS*, 1-5.

Characteristics and Seasonal Abundance of Seagrass at Pidakon Coast Pacitan, East Java, Indonesia

ORIGINALITY REPORT

15%

SIMILARITY INDEX

8%

INTERNET SOURCES

8%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|----|
| 1 | Lina Mtwana Nordlund, Emma L. Jackson, Masahiro Nakaoka, Jimena Samper-Villarreal, Pedro Beca-Carretero, Joel C. Creed. "Seagrass ecosystem services – What's next?", Marine Pollution Bulletin, 2017
Publication | 1% |
| 2 | Submitted to Townsville Grammar School
Student Paper | 1% |
| 3 | www.marine.uq.edu.au
Internet Source | 1% |
| 4 | Submitted to National Institute of Education
Student Paper | 1% |
| 5 | www.oceandocs.org
Internet Source | 1% |
| 6 | T. Thangaradjou, J.R. Bhatt. "Status of seagrass ecosystems in India", Ocean & Coastal Management, 2017
Publication | 1% |

7	R. Purvaja, R.S. Robin, D. Ganguly, G. Hariharan, G. Singh, R. Raghuraman, R. Ramesh. "Seagrass meadows as proxy for assessment of ecosystem health", Ocean & Coastal Management, 2018 Publication	1%
8	www.tsra.gov.au Internet Source	1%
9	Submitted to Deakin University Student Paper	1%
10	www.redalyc.org Internet Source	1%
11	Campbell, S.J.. "Photosynthetic responses of seven tropical seagrasses to elevated seawater temperature", Journal of Experimental Marine Biology and Ecology, 20060321 Publication	1%
12	S. Kupschus. "Associations between fish assemblages and environmental factors in nearshore habitats of a subtropical estuary", Journal of Fish Biology, 5/2001 Publication	1%
13	Submitted to University of Cape Town Student Paper	1%
14	Unsworth, Richard K.F., Catherine J. Collier,	1%

Michelle Waycott, Len J. Mckenzie, and Leanne C. Cullen-Unsworth. "A framework for the resilience of seagrass ecosystems", Marine Pollution Bulletin, 2015.

Publication

15

L.G. Egea, R. Jiménez-Ramos, J.J. Vergara, I. Hernández, F.G. Brun. "Interactive effect of temperature, acidification and ammonium enrichment on the seagrass *Cymodocea nodosa*", Marine Pollution Bulletin, 2018

Publication

1 %

16

www.aab.bioflux.com.ro

Internet Source

1 %

17

rrrc.com.au

Internet Source

<1 %

18

Submitted to Bath Spa University College

Student Paper

<1 %

19

pir.sa.gov.au

Internet Source

<1 %

20

"Encyclopedia of Modern Coral Reefs", Springer Nature, 2011

Publication

<1 %

21

www.gbrmpa.gov.au

Internet Source

<1 %

22

Kathryn M. Chartrand, Milán Szabó, Sutinee

<1 %

Sinutok, Michael A. Rasheed, Peter J. Ralph.
"Living at the margins – The response of deep-
water seagrasses to light and temperature
renders them susceptible to acute impacts",
Marine Environmental Research, 2018

Publication

23

journal.unnes.ac.id

Internet Source

<1%

Exclude quotes On

Exclude matches < 10 words

Exclude bibliography On